



ARL-TR-8204 • Nov 2017



Base Stability of Aminocyclopropeniums

by MyVan Baranoski, Frederick L Beyer, Robert Lambeth, and
Nicole Zander

Approved for public release; distribution is unlimited.

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.



Base Stability of Aminocyclopropeniums

**by MyVan Baranoski, Frederick L Beyer, Robert Lambeth, and
Nicole Zander**

Weapons and Materials Research Directorate, ARL

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) November 2017		2. REPORT TYPE Technical Report		3. DATES COVERED (From - To) 01 September 2016–30 September 2017	
4. TITLE AND SUBTITLE Base Stability of Aminocyclopropeniums				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) MyVan Baranoski, Frederick L Beyer, Robert Lambeth, and Nicole Zander				5d. PROJECT NUMBER H94	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Research Laboratory Weapons and Materials Research Directorate (ATTN: RDRL-WMM-G) 2800 Powder Mill Road Adelphi, MD 20783-1138				8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TR-8204	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <p>Cyclopropenium cations with differing amino-functional groups were synthesized and tested for their utility in anion exchange membranes for alkaline fuel cells. A series of aminocyclopropeniums were synthesized and their base stability probed in situ using time-resolved proton nuclear magnetic resonance (¹H NMR) under conditions designed to mimic those encountered in alkaline fuel cells to test their utility in anion exchange membranes. While the aminocyclopropeniums showed poor base stability, the cyclopropenium cation may have beneficial properties in neutral pH applications such as antimicrobial fabrics and coatings.</p>					
15. SUBJECT TERMS cyclopropenium, anion exchange membrane, alkaline stability, aminocyclopropenium, cation					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 16	19a. NAME OF RESPONSIBLE PERSON MyVan Baranoski
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) (410) 306-0293

Contents

List of Figures	iv
1. Introduction	1
2. Methods and Procedures	1
2.1 General Syntheses of Tris(dialkylamino)cyclopropenium Chloride	2
2.2 Synthesis of Tris(diethylamino)cyclopropenium Chloride #1	2
2.3 Synthesis of Tris(dipropylamino)cyclopropenium Chloride #2	2
2.4 Synthesis of Tris(dibutylamino)cyclopropenium Chloride #3	3
2.5 Synthesis of Diethylaminobis(diisopropylamino)cyclopropenium Chloride #4	3
2.6 Synthesis of Diethylaminobis(dicyclohexylamino)cyclopropenium Chloride #5	3
3. Results and Discussion	4
4. Conclusions	6
5. References	8
List of Symbols, Abbreviations, and Acronyms	9
Distribution List	10

List of Figures

Fig. 1	Selected aminocyclopropenium cations for alkaline stability	4
Fig. 2	Time resolved ^1H NMR of tris(diethylamino)cyclopropenium cation in 1M KOH/ CD_3OH at 80 $^\circ\text{C}$	5
Fig. 3	Fraction of aminocyclopropenium cations remaining (left) in 1M KOH/ CD_3OH at 80 $^\circ\text{C}$, as determined from ^1H NMR data.....	5
Fig. 4	Alkaline hydrolysis of aminocyclopropeniums	6

1. Introduction

Polymers containing cationic functional groups have the potential to be used in a wide range of applications, including antimicrobial coatings and fabrics, electrolyzers, anion exchange membranes in alkaline fuel cells, and redox flow batteries.^{1,2} These applications present challenging materials science problems because of the orthogonal properties that are typically required. For example, in alkaline fuel cells, high affinity for hydroxide (OH^-) is required, although hydroxide causes degradation of the cation needed for that affinity.

Most of the work in cation-containing polymers focused on a small number of functional groups including ammonium, phosphonium, imidazolium, and pyridinium ions, which are generally application specific with limited ability to tune physical properties.³ Their successful use in alkaline fuel cells has been hindered due to the highly nucleophilic nature of the hydroxide anion, which is able to attack the covalently attached cation through multiple reaction pathways, rendering the material neutral.^{2,4} In addition, compared to ionic liquids, ion mobility in solid polyelectrolytes is significantly reduced, with the movement of free anions limited to the interstitial space between the polymer chains.

A key issue with anion exchange membranes is the cation lifetime in a high pH environment ($>\text{pH } 13$). The cyclopropenium ion is an aromatic and unusually electron-rich cation that has improved dispersion of charge relative to many other cations.^{5,6} A new class of polyelectrolytes was developed from cyclopropenium ions by Killops et al.⁵ These materials are predicted to have high thermodynamic stability,⁷ and Killops' data suggest this to be the case.

In this work, cyclopropenium cations with differing amino-functional groups were synthesized and tested for their utility in anion exchange membranes for alkaline fuel cells. To determine cyclopropenium ion alkaline stability, a series of aminocyclopropeniums were synthesized and their base stability probed in situ using time-resolved proton nuclear magnetic resonance (^1H NMR) under conditions designed to mimic those encountered in alkaline fuel cells to test their utility in anion exchange membranes.

2. Methods and Procedures

All operations were performed using standard Schlenk techniques under a nitrogen atmosphere to reduce exposure to water. The ^1H NMR spectra were collected using a Bruker 600 MHz Avance instrument in deuterated chloroform (CDCl_3), referenced to residual solvent peaks.

Pentachlorocyclopropane was prepared from sodium trichloroacetate and trichloroethylene.⁸ The amines were used as obtained commercially. All reagents were obtained from commercial sources and used as received unless otherwise noted.

2.1 General Syntheses of Tris(dialkylamino)cyclopropenium Chloride⁹

Pentachlorocyclopropane (1 equivalent) was added to dry dichloromethane (CH_2Cl_2) (approximately 0.01 M). This was stirred at 0 °C under a nitrogen atmosphere. Eight equivalents of amine were slowly added. Following the complete addition of the amine, the solution was stirred at 0 °C for a further 4 h, and allowed to come to ambient temperature and stirred overnight (16 h) before heating to reflux (65 °C) for 5 h. The solution, usually yellow/orange in color, was then placed under vacuum to remove the solvent.

2.2 Synthesis of Tris(diethylamino)cyclopropenium Chloride #1

Pentachlorocyclopropane ($\text{C}_3\text{Cl}_5\text{H}$, 0.51 g, 2.33 mmol) and diethylamine (HNEt_2 , 1.34 g, 18.7 mmol) were combined using the method listed under General Syntheses. Dichloromethane was removed in vacuo and 20 mL of acetone was added to precipitate diethylammonium chloride, which was then filtered from the solution comprising the triaminocyclopropenium salt. Acetone was removed in vacuo and the resulting solid was dissolved in 20 mL of CH_2Cl_2 . The organic solution was washed with dilute sodium hydroxide (NaOH) (15 mL) followed by hydrochloric acid (HCl) (15 mL) and then water (H_2O) (3×15 mL). The organic solution was dried over sodium sulfate (Na_2SO_4). The organic solvent was removed under vacuum to give an orange liquid (0.59 g, 86% yield) that slowly solidified under high vacuum overnight. ^1H NMR (CDCl_3 , 600 MHz): δ 3.42 (q, 12H, NCH_2), 1.27 (t, 18H, NCH_2CH_3).

2.3 Synthesis of Tris(dipropylamino)cyclopropenium Chloride #2

$\text{C}_3\text{Cl}_5\text{H}$ (0.51 g, 2.33 mmol) and dipropylamine (HNPr_2 , 1.92 g, 18.7 mmol) were combined using the method listed under General Syntheses. Dichloromethane was removed in vacuo and 20 mL of acetone was added to precipitate dipropylammonium chloride, which was then filtered from the solution comprising the triaminocyclopropenium salt. Acetone was removed in vacuo and the resulting solid was dissolved in 20 mL of CH_2Cl_2 . The organic solution was washed with H_2O (3×15 mL). The organic solution was dried over Na_2SO_4 . The organic solvent

was removed under vacuum to give an orange liquid (0.64 g, 75% yield) that slowly solidified under high vacuum overnight. ^1H NMR (CDCl_3 , 600 MHz): δ 3.26 (t, 12H, NCH_2), 1.64 (m, 12H, NCH_2CH_2), 0.92 (t, 18H, $\text{NCH}_2\text{CH}_2\text{CH}_3$).

2.4 Synthesis of Tris(dibutylamino)cyclopropenium Chloride #3

$\text{C}_3\text{Cl}_5\text{H}$ (0.51g, 2.33 mmol) and dibutylamine (HNBu_2 , 2.4 g, 18.7 mmol) were combined using the method listed under General Syntheses. Dichloromethane was removed in vacuo and 20 mL of acetone was added to precipitate dipropylammonium chloride, which was then filtered from the solution comprising the triaminocyclopropenium salt. Acetone was removed in vacuo and the resulting solid was dissolved in 20 mL of CH_2Cl_2 . The organic solution was washed with H_2O (3×15 mL). The organic solution was dried over Na_2SO_4 and the solvent was removed under vacuum. The remaining liquid was washed with diethyl ether (4×15 mL) to remove residual amine (0.93 g, 86% yield). ^1H NMR (CDCl_3 , 600 MHz): δ 3.29 (t, 12H, NCH_2), 1.59 (m, 12H, NCH_2CH_2), 1.31 (m, 12H, $\text{NCH}_2\text{CH}_2\text{CH}_3$), 0.94 (t, 18H, CH_3).

2.5 Synthesis of Diethylaminobis(diisopropylamino)cyclopropenium Chloride #4

$\text{C}_3\text{Cl}_5\text{H}$ (0.5g, 2.33 mmol) was added to dry CH_2Cl_2 . This was stirred at 0 °C under a nitrogen atmosphere. Diisopropylamine ($\text{HN}(\text{iPr})_2$, 1.4 g, 14.0 mmol) was added dropwise using a syringe pump and the solution was allowed to return to room temperature and stirred for approximately 36 h. HNEt_2 (0.34 g, 4.65 mmol) was added dropwise and the reaction was stirred for an additional 5 days. Approximately 3 mL of concentrated HCl was added to precipitate excess amine, which was then filtered from the solution comprising the aminocyclopropenium salt. The solution was washed with 1M HCl ($4 \text{ mL} \times 5$) to remove any residual amine. The solution was dried over Na_2SO_4 and the solvent was removed under vacuum (0.59 g, 74% yield). ^1H NMR (CDCl_3 , 600 MHz): δ 3.86 (m, 4H, $\text{N}(\text{iPr})\text{H}$), 3.56 (q, 4H, NCHCH_3), 1.36 (d, 24H, $\text{N}(\text{iPr})\text{H}$), 1.27 (t, 6H, NCH_2CH_3).

2.6 Synthesis of Diethylaminobis(dicyclohexylamino)cyclopropenium Chloride #5

$\text{C}_3\text{Cl}_5\text{H}$ (0.5g, 2.33 mmol) was added to dry CH_2Cl_2 . This was stirred at 0 °C under a nitrogen atmosphere. HNCy_2 (2.54 g, 14.0 mmol) was added dropwise using a syringe pump and the solution was allowed to return to room temperature and stirred for approximately 36 h. HNEt_2 (0.34 g, 4.65 mmol) was added dropwise and the reaction was stirred for an additional 12 h. Approximately 3 mL of concentrated

HCl was added to precipitate excess amine, which was then filtered from the solution comprising the aminocyclopropenium salt. The solution was dried over Na₂SO₄ and the solvent was removed under vacuum. The crude product was recrystallized in hexanes (0.81 g, 69% yield). ¹H NMR (CDCl₃, 600 MHz): δ 3.57 (q, 4H, NCH₂CH₃), 3.33 (m, 4H, NCyH), 1.05-1.95 (m, 46H, CyH, NCH₂CH₃).

The base stability of the aminocyclopropeniums was probed in situ using time resolved ¹H NMR under conditions designed to mimic those encountered in alkaline fuel cells. A 1M potassium hydroxide (KOH) stock solution was prepared by dissolving KOH (323.6 mg, 5.77 mmol) and 1,4-dioxane (12.7 mg, 0.144 mmol) in 5.77 mL of deuterated methanol (CD₃OH). Dioxane is used as an internal standard; 0.0375 mmol of aminocyclopropenium was dissolved in the stock solution (0.75 mL) and transferred into a nuclear magnetic resonance (NMR) tube. The tube was then placed in the spectrometer, and heated to 80 °C. The resonances associated with the alkyl substituents were then recorded over time. Integration of a selected signal in the model compound relative to the signal related to dioxane provided the quantity of model compound.

3. Results and Discussion

Figure 1 illustrates the 5 cyclopropenium cations with different amino substituents that were explored in this study. Substituents were selected to determine if structural factors affect the base stability of aminocyclopropeniums.




Fig. 1 Selected aminocyclopropenium cations for alkaline stability

The base stability was determined by dissolving the aminocyclopropenium in a 1M KOH stock solution and probing in situ using time resolved ^1H NMR for 14 h at 80 °C. A typical example of the NMR data is shown in Fig 2. The selected signal of the model compound at 3.45 ppm integrated relative to the signal related to dioxane at 3.66 ppm provided the quantity of the compound.




Fig. 2 Time resolved ^1H NMR of tris(diethylamino)cyclopropenium cation in 1M KOH/ CD_3OH at 80 °C

The results of the study are illustrated in Fig. 3. The aminocyclopropeniums show 5%–15% degradation within 30 min. After 6.5 h, there is less than 50% cation remaining.




Fig. 3 Fraction of aminocyclopropenium cations remaining (left) in 1M KOH/ CD_3OH at 80 °C, as determined from ^1H NMR data

The aminocyclopropeniums with simple alkyl and bulky substituents showed poor base stability, with only 5%–20% of cation remaining after 14 h. In comparison, a study of a tetrakis(dialkylamino)phosphonium cation showed no degradation over 20 days at 80 °C.¹⁰ To determine if poor cation base stability translates to polymers with poor stability, a cyclopropenium polymer with isopropylamine substituents was also evaluated using the same time-resolved NMR method and found to also quickly degrade under the same conditions, suggesting that cyclopropenium is not stable under these conditions.

These results could be explained by the findings reported by Yoshida et al.¹¹ where aminocyclopropeniums undergo hydrolysis in alkaline conditions at room temperature to give a diaminocyclopropenone and an acrylamide derivative (Fig. 4).




Fig. 4 Alkaline hydrolysis of aminocyclopropeniums

Yoshida et al. further explained that the distribution of the products appeared to be dependent on the basicity of the amino groups attached to the cyclopropenium ring, the alkali used, and its concentration. As an example, 1,2,3-tris-(N-methylanilino)cyclopropenium perchlorate in methanol was stirred at room temperature with 0.3% aq KOH for 40 h. This yielded the cyclopropenone at 22% and the acrylamide at 65%. Alkaline hydrolysis is the most likely degradation route for the aminocyclopropeniums examined in this study.

4. Conclusions

This report details the synthesis of various aminocyclopropenium salts with differing alkyl substituents and their base stability. The aminocyclopropeniums showed poor base stability, showing 5%–15% degradation within 30 min, less than 50% cation remaining after 6.5 h, and only 5%–20% of cation remaining after 14 h. Other cations had little to no degradation under similar conditions over longer

time periods. These results are consistent with literature findings where aminocyclopropeniums undergo alkaline hydrolysis at room temperature. While base stability hinders their utility as anion exchange membranes in alkaline fuel cells, the cyclopropenium cation may have beneficial properties in neutral pH applications such as antimicrobial fabrics and coatings.

5. References

1. Carmona-Ribeiro AM, de Melo Carrasco LD. Cationic antimicrobial polymers and their assemblies. *Int J Mol Sci.* 2013;14(5):9906–9946.
2. Varcoe JR, Atanassov P, Dekel DR, Herring AM, Hickner MA, Kohl PA, Kucernak AR, Mustain WE, Nijmeijer K, Scott K, et al. Anion-exchange membranes in electrochemical energy systems. *Energy Environ Sci.* 2014;7(10): 3135–3191.
3. Yuan J, Antonietti M. Poly(ionic liquid)s: polymers expanding classical property profiles. *Polymer.* 2011;52(7):1469–1482.
4. Yuan J, Mecerreyes D, Antonietti M. Poly(ionic liquid)s: an update. *Prog Polym Sci.* 2013;38:1009–1036.
5. Killops KL, Brucks SD, Rutkowski KL, Freyer JL, Jiang Y, Valdes ER, Campos LM. Synthesis of robust surface-charged nanoparticles based on cyclopropenium ions. *Macromolecules.* 2015;48(8):2519–2525.
6. Jiang Y, Freyer JL, Cotanda P, Brucks SD, Killops KL, Bandar JS, Torsitano C, Balsara NP, Lambert TH, Campos LM. The evolution of cyclopropenium ions into functional polyelectrolytes. *Nat Commun* 2015;6(5950).
7. Jemmis ED, Subramanian G, Kos AJ, Schleyer PVR. The remarkably stabilized trithiocyclopropenium ion, $C_3Li_3^+$, and its relatives. *J Am Chem Soc.* 1997;119:9504–9512.
8. Tobey SW, West R. Pentachlorocyclopropane. *J Am Chem Soc.* 1966;88(11):2478–2481.
9. Curnow OJ, MacFarlane DR, Walst KJ. Triaminocyclopropenium salts as ionic liquids. *Chem Commun.* 2011;47:10248–10250.
10. Noonan KJT, Hugar KM, Kostalik HA 4th, Lobkovsky EB, Abruna HD, Coates GW. Phosphonium-functionalized polyethylene: a new class of base-stable alkaline anion exchange membranes. *J Am Chem Soc.* 2012;134:18161–18164.
11. Yoshida Z, Konishi H, Tawara Y, Nishikawa K, Ogoshi H. Novel alkaline hydrolysis of triaminocyclopropenium ion. New route to diaminocyclopropenone and diaminocyclopropenethione. *Tetrahedron Letters.* 1973;14(28):2619–2622.

List of Symbols, Abbreviations, and Acronyms

^1H NMR	proton nuclear magnetic resonance
ARL	US Army Research Laboratory
$\text{C}_3\text{Cl}_5\text{H}$	pentachlorocyclopropane
CD_3OH	deuterated methanol
CDCl_3	deuterated chloroform
CH_2Cl_2	dichloromethane
H_2O	water
HCl	hydrochloric acid
$\text{HN}(\text{iPr})$	diisopropylamine
HNPr_2	dipropylamine
HNBu_2	dibutylamine
HNEt_2	diethylamine
KOH	potassium hydroxide
mmol	millimole
Na_2SO_4	sodium sulfate
NaOH	sodium hydroxide
NCH_2CH_3	triethylamine
NMR	nuclear magnetic resonance
OH^-	hydroxide
pH	potential of hydrogen

1 DEFENSE TECHNICAL
(PDF) INFORMATION CTR
DTIC OCA

2 DIR ARL
(PDF) IMAL HRA
RECORDS MGMT
RDRL DCL
TECH LIB

1 GOVT PRINTG OFC
(PDF) A MALHOTRA

4 DIR ARL
(PDF) RDRL WMM G
M BARANOSKI
F BEYER
R LAMBETH
N ZANDER